

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 1 182 257 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
27.02.2002 Bulletin 2002/09

(51) Int Cl<sup>7</sup>: **C12N 15/53, C12N 9/02,  
C12N 9/04, C12N 15/82,  
A01H 5/00**

(21) Application number: **00308002.5**

(22) Date of filing: **14.09.2000**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

- Yi, Hankul  
Kwangju-city (KR)
- SHIN, Byongchul  
Kwangju-city (KR)

(30) Priority: **14.08.2000 US 638715**

(74) Representative: **Nash, David Allan et al  
Hasseltine Lake & Co., Imperial House, 15-19  
Kingsway  
London WC2B 6UD (GB)**

(71) Applicant: **Korea Kumho Petrochemical Co. Ltd.  
Chongno-Gu, Seoul (KR)**

Remarks:

The sequence listing, which is published as annex to the application documents, was filed after the date of filing. The applicant has declared that it does not include matter which goes beyond the content of the application as filed.

(72) Inventors:  
• **CHOI, Giltso**  
Kwangju-city (KR)  
• **JOHNSON, Eric T.**  
Pullman - WA 99163 (US)

(54) **Nucleic acids encoding dihydroflavonol 4-reductase with altered specificity and uses thereof**

(57) The invention includes a modified dihydroflavonol 4-reductase (DFR) nucleic acids encoding a modified DFR that has altered amino acid sequences at the substrate specificity determining region. This modified DFR is characterized by its ability to reduce dihydrokaempferol (DHK) preferentially over dihydroquercetin (DHQ), and dihydromyricetin (DHM). The invention also includes plants having at least one cell ex-

pressing the modified DFR. Such plants are characterized by an increased content of pelargonidin-based pigments. The invention also includes vectors comprising at least a fragment of the nucleic acids encoding said modified DFR. The invention also includes methods using such vectors for producing plants having an increased content of pelargonidin-based pigments.

EP 1 182 257 A1

## Description

## BACKGROUND OF THE INVENTION

## Technical field of the invention

[0001] The present invention relates to modified DFR nucleic acids and encoding the modified DFR that preferentially utilize DHK as a substrate and their uses for genetically altering plants to increase the content of pelargonidin-based pigments in the plants.

## Description of the Prior art

[0002] Anthocyanins are classes of pigments that determine flower color and plant pigmentation in angiosperm plants. Among anthocyanins, pelargonidin-based pigments confer brick-red/orange color to plants, while cyanidin- and delphinidin-based pigments confer red and violet color each (Holton, et al. Plant Cell 7:1071-1083 (1995); Tanaka, et al. Plant Cell Physiol. 39:1119-1126 (1998)). Different ratio of these pigments confers a wide range of flower color. Many anthocyanin biosynthetic genes have been identified. One of key enzyme in the biosynthetic pathway is dihydroflavonol 4-reductase (DFR). The enzyme converts dihydroflavonols (dihydrokaempferol (DHK), dihydroquercetin (DHQ), and dihydromyricetin (DHM)) to leucocyanidins. The leucocyanidins are subsequently converted to anthocyanins by other enzymes. The conversion of DHK to DHQ and DHM are catalyzed by flavonoid 3'-hydroxylase (F3'H) and flavonoid 3',5'-hydroxylase (F3',5'H). Since DFRs in most plants can convert all three dihydroflavonols to leucocyanidins, the ratio of three classes of anthocyanin pigments are mainly determined by the activity of F3'H and F3',5'H (Holton, et al. Plant Cell 7:1071-1083 (1995)).

[0003] Since pelargonidin-based pigments confer the orange color to flowers, the F3'H and F3',5'H activities must be absent for a plant to have orange colored flowers (U.S. patent 5410096). In many plant species, F3'H and F3',5'H are encoded by multiple genes, thus the mutant lines that lack F3'H and F3',5'H are not easily found. This partially accounts for the rarity of orange-colored flowers in some plant species. Inability to reduce DHK to leucocyanidin by DFR in some species can also cause the lack of orange-colored flower. For example, DFRs from *Petunia* and *Cymbidium* convert DHK to its leucocyanidin very inefficiently, thus these species do not accumulate large ratio of pelargonidin-based anthocyanins even if F3'H and F3',5'H are absent (Gerats, et al. Planta 155:364-368 (1982); Johnson, et al. Plant J. 19:81-85 (1999)). An orange-colored *Petunia* was engineered by introducing a maize DFR to a special mutant line of *Petunia* that lacks F3'H and F3',5'H (Meyer, et al. Nature 330:677-678 (1987)). Since the maize DFR can convert all three dihydroflavonols to their leucocyanidins, such a mutant line that accumulates DHK was necessary for the development of orange-colored *Petunia*. The necessity of the special mutant line can be circumvented by using a DFR that utilizes DHK preferentially over DHQ and DHM.

[0004] Using chimeric DFRs between *Petunia* and *Gerbera* DFRs, we identified a region that determines the substrate specificity of DFR. By altering an amino acid in the region, we developed a DHK-specific DFR that converts DHK preferentially over DHQ and DHM. When expressed in plants, the DHK-specific DFR increases the pelargonidin-based pigments regardless of F3'H activity.

## SUMMARY OF THE INVENTION

[0005] Accordingly, the object of this invention is to provide substrate-specific DFRs which have altered amino acid sequences at the substrate specificity determining region.

[0006] It is an also object herein to provide a DHK-specific DFR and nucleic acids encoding the DHK-specific DFR.

[0007] Still further, it is an object herein to provide transgenic plants expressing the DHK-specific DFR which confers a phenotype characterized by the increased content of pelargonidin-based pigments in the plants.

[0008] In accordance with the objects, the invention includes the modified DFRs and nucleic acids encoding the modified DFRs which have altered amino acid sequences at the substrate specificity determining region. The properties of modified DFRs are characterized by their abilities to reduce one substrate preferentially among DHK, DHQ, and DHM.

[0009] The invention also includes a modified DFR that reduces DHK preferentially over DHQ and DHM.

[0010] The invention also includes plants having at least one cell transformed with a vector comprising at least a portion of the modified DFR nucleic acids. Such plants have a phenotype characterized by the increased content of pelargonidin-based pigments.

[0011] The invention also includes vectors capable of transforming a plant cell to increase the content of pelargonidin-based pigments.

[0012] The invention also includes methods for producing plants having the increased content of pelargonidin-based pigments. The methods includes steps of transforming plant cells with vectors containing the modified DFR gene;

regenerating plants from the transformed cells and selecting the plant having the increased content of pelargonidin-based-pigments.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0013] Figure 1A is a schematic diagram showing three chimeric *DFRs*. Black bars indicate sequences from a *Gerbera DFR* and gray bars indicate sequences from a *Petunia DFR*. Numbers are junctional amino acid positions from the translation start site of the *Gerbera DFR*. C.1, C.2, C.3 are the name of three different chimeric *DFRs*.

[0014] Figure 1B shows representative flowers of transgenic *Petunia* expressing chimeric *DFRs* or control *DFR*. Ger indicate the transgenic flower expressing *Gerbera DFR* and C.1, C.2, and C.3 indicate Chimera 1, Chimera 2, and Chimera 3 each. RLO1 line has a functional *Petunia DFR* gene. The C.1 and RLO1 bore similar pink colored flowers while others bore brick-red colored flower. The transgenic W80 flower expressing C.1 has pink color, while transgenic W80 flowers expressing C.2 and C.3 have orange/brick-red color. The orange/brick-red color can be also observed in the transgenic *Petunia* flowers expressing the native *Gerbera DFR*.

[0015] Figure 1C shows the TLC analysis data of pigments produced in transgenic *Petunia* flowers next to standard pigments (pelargonidin (Pg), cyanidin (Cy), and delphinidin (Dp)). The transgenic flowers expressing C.1 has mainly cyanidin- and delphinidin-based pigments, while the flowers expressing C.2 and C.3 have mainly pelargonidin-based pigments in addition to small amount of cyanidin- and delphinidin-based pigments.

[0016] Figure 2 shows the amino acid sequence of *Gerbera DFR* aligned with other representative *DFR* sequences. The ClustalW program was used to align multiple amino acid sequences (Thomson, et al. Nucl. Acids Res. 22: 4673-4680 (1994)). The substrate specificity determining region is boxed and the 134<sup>th</sup> amino acid residue of *Gerbera DFR* and corresponding amino acid residues of *DFRs* from a few representative species are bold typed.

[0017] Figure 3A shows site-directed mutagenesis of substrate specificity determining region. The sequence corresponds to the substrate specificity determining region of *Gerbera DFR*. Arrows and letters indicates amino acids that were changed to.

[0018] Figure 3B shows flowers of transgenic *Petunia* expressing mutated *Gerbera DFR* gene. Ger indicates the wild type *Gerbera DFR* and T132V indicates the mutated *DFR* that has valine instead of threonine at the 132<sup>th</sup> position of *Gerbera DFR*. Names of other mutated *DFRs* followed the same notation rule. All transgenic lines except N134L and E145L have the same brick red colored flower. The N134L bore slightly different colored flowers and E145L bore white flowers.

[0019] Figure 3C shows a TLC analysis of pigments produced in the transgenic *Petunia* flowers. As expected, the E145L did not accumulate any anthocyanin. The N134L accumulated mostly pelargonidin while other mutated *DFR* and wild type *Gerbera DFR* accumulated significant amount of cyanidin and delphinidin in addition to pelargonidin.

[0020] Figure 4A shows the development of a *DFR* that display the altered substrate specificity. WR and WV indicate *Petunia* lines that are *dhfr<sup>-</sup>*, but *F3'H<sup>+</sup>* (WR) or *F3'5'H<sup>+</sup>* (WV). The mark - indicates no *DFR* gene, *DFR<sup>N134L</sup>* indicates *DFR* that has leucine instead of asparagine at the 134<sup>th</sup> position of *Gerbera DFR*, and *DFR<sup>WT</sup>* indicates the wild type *Gerbera DFR*. The flower located in the cross section indicate the WR or WV transgenic flowers expressing *DFR<sup>N134L</sup>* or *DFR<sup>WT</sup>*.

[0021] Figure 4B shows a TLC analysis of pigments produced in the transgenic lines. Pg, Cy, and Dp indicate pelargonidin, cyanidin, and delphinidin. The WR and WV lines expressing wild type *DFR* accumulated cyanidin and delphinidin each. The WR line expressing *DFR<sup>N134L</sup>* accumulated pelargonidin and cyanidin, while the WV line expressing *DFR<sup>N134L</sup>* did not accumulate any pigment other than background level of delphinidin.

## DETAILED DESCRIPTION OF THE INVENTION

[0022] In accordance with the present invention, the substrate specificity determining region was identified by determining the abilities of three chimeric *DFRs* to catalyze the reduction of DHK in the transgenic *Petunia* lines. In order to identify the region of *DFR* that determines its substrate specificity, we constructed chimeric *DFR* genes using cDNA sequences of *Petunia* and *Gerbera*. Though these two *DFRs* have high similarity at the amino acid level, *Gerbera DFR* is able to catalyze dihydrokaempferol (DHK) while *Petunia DFR* cannot (Elomaa et al. Mol. Gen. Genet. 248:649-656 (1995)). We built three different chimeric genes using regions of high homology as common PCR primer sites (Fig1A). The chimeric genes were transformed into a white flowered *Petunia* mutant (W80) that lacks *DFR* activity and accumulates primarily DHK but with appreciable amounts of dihydroquercetin (DHQ) and dihydromyricetin (DHM) (Huits et al., 1994). Chimera 1 produced pink flowers while Chimeras 2 and 3 bore orange-pink flowers (Fig. 1B). The hue of Chimera 1 flowers is very similar to the inbred *Petunia* mutant RLO1, which has functional *DFR* activity and accumulates DHK. Thin layer chromatography (TLC) determined that Chimera 1 produced mainly cyanidin and delphinidin (Fig. 1b). Chimeras 2 and 3 primarily produced pelargonidin (Fig. 1C), which is the downstream product of *DFR* reduction of DHK. These results indicated that the region of *DFR* conferring the ability to reduce DHK was between Chimeras 1

and 2. The identified region (approx. 40 amino acids) is highly variable in DFRs from different plant species. By excluding the completely conserved amino acid sequences at the borders, the identified region is narrowed down to 26 amino acids. Hereinafter, this region is referred as substrate specificity determining region. An example of the substrate specificity determining region in a few representative DFRs is shown in Figure 2.

[0023] The invention provides the modified *DFR* nucleic acids and encoded DFRs that have altered amino acid sequences at the substrate specificity determining region. Such DFRs have properties characterized by the altered substrate specificity. Hereinafter, DFRs that catalyze the reduction of one substrate preferentially over other two substrates are referred as substrate-specific DFRs. In the preferred embodiments, the invention provides the modified DFR that has altered amino acid at 134<sup>th</sup> amino acid residue of *Gerbera* DFR or the corresponding amino acid residues of DFRs from other species. Such DFRs have properties characterized by catalyzing the reduction of DHK preferentially over DHQ and DHM. Hereinafter, DFRs that catalyze the reduction of DHK preferentially over DHQ and DHM are referred as DHK-specific DFRs. The 134<sup>th</sup> amino acid residue of *Gerbera* DFR and corresponding amino acid residues of DFRs from a few representative species are shown in Figure 2.

[0024] In accordance with the present invention, a DHK-specific DFR was developed by replacing asparagine at 134<sup>th</sup> amino acid residue of *Gerbera* DFR to leucine. The expression of the DHK-specific DFR in W80 *Petunia* line, which accumulates large amount of DHK in addition to appreciable amount of DHQ and DHM, caused the production of only pelargonidin. The expression of native *Gerbera* DFR in the same *Petunia* line caused the production of appreciable amounts of cyanidin and delphinidin in addition to pelargonidin (Figure 3). Since the W80 *Petunia* line we transformed accumulates mainly DHK with small amount of DHQ and DHM, it was not clear if the N134L mutant DFR completely lost the capability of reducing DHQ and DHM. To investigate if the N134L mutant DFR produces only pelargonidin in the presence of fully active flavonoid-3'-hydroxylase (F3'H) or flavonoid-3',5'-hydroxylase (F3'5'H), we crossed our N134L transformant with *Petunia* lines that are either *dfr<sup>-</sup>/F3'H<sup>+</sup>* (WR line) or *dfr<sup>-</sup>/F3'5'H<sup>+</sup>* (WV line). As shown in figure 4A, both WR and WV lines bore white flowers as expected. When these lines were crossed with the N134L transformants, the WR line expressing the mutant DFR (WR/DFR<sup>N134L</sup>) had orange colored flowers while the WR expressing wild type *Gerbera* DFR (WR/DFR<sup>WT</sup>) had red colored flowers. Unlike the WR lines, the WV lines expressing the mutant DFR (WV/DFR<sup>N134L</sup>) bore white flowers while WV lines expressing the wild type DFR (WV/DFR<sup>WT</sup>) had violet colored flowers. To determine the pigments produced in these crossed lines, we performed TLC analysis. Figure 4B shows that the WR/DFR<sup>N134L</sup> accumulated a large amount of pelargonidin while WR/DFR<sup>WT</sup> mainly accumulated cyanidin. In the white flowered WV/DFR<sup>N134L</sup>, no appreciable amounts of anthocyanidins accumulated other than a background level of delphinidin. In contrast to WV/DFR<sup>N134L</sup>, the WV/DFR<sup>WT</sup> accumulated mainly delphinidin. The data indicate that the N134L mutant DFR preferentially utilizes DHK as a substrate over DHQ and cannot reduce DHM. The substrate preference of the N134L mutant DFR is somewhat opposite to that of *Petunia* DFR which prefer DHM over DHQ and cannot use DHK (Forkmann and Ruhnau, 1987). The results indicates that the DHK-specific DFR can increase the pelargonidin-based pigments in plants regardless of the presence of F3'H activity.

[0025] The invention also provides plants having cells transformed with vectors comprising at least a portion of the substrate-specific *DFR* nucleic acids. Such plants have phenotypes characterized by the increased content of anthocyanins specified by the substrate specific *DFRs*. In the preferred embodiments, the invention provides plants having cells transformed with vectors comprising at least a portion of the DHK-specific *DFR* nucleic acids. Such plants have phenotypes characterized by the increased content of pelargonidin-based pigments. Plants that can be used to practice the invention include plants within the Division of Magnoliophyta, i.e. the angiosperms include the dicotyledons and the monocotyledons. Particularly preferred Orders of angiosperms according to "Plant Systematics", S.B. Jones, Jr. and A.E. Luchsinger include Magnoliales, Laurales, Aristolochiales, Nymphaeales, Ranunculales, Caryophyllales, Malvales, Violales, Capparales, Ericales, Primulales, Rosales, Fabales, Myrtales, Cornales, Rhamiales, Sapindales, Geraniales, Apiales, Gentianales, Solanales, Lamiales, Scrophulariales, Campanulales, Rubiales, Dipsacales, Asterales, Hydrocharitales, Arales, Cyperales, Liliales, and Orchidales. Particularly preferred plants include orchid, iris, campanula, gentiana, phlox, cyclamen, eustoma, crocus, delphinium, ageratum, chrysanthemum, *Petunia*, cactus, limonium, astilbe, carnation, *Gerbera*, brassica, impatiens, geranium, dahlia, sunflower, dianthus, gloxinia, caledula, bellis, ranunculus, aster, tagetes, salvia, hibiscus, cirsium, godetia, catharanthus, alyssum, lupinus, portulaca, drosera, tulip, lily, narcissus, freesia, anemone, gladiolus, caladium, archimenes, achillea, agapanthus, aethiones, allium, alstroemeria, amaryllis, anagallis, androsace, anemone, antirrhinum, aquilegia, ameria, asperula, begonia, browallia, callistephus, camellia, ceanothus, chionodoxa, cistus, clarkia, clematis, colchicum, consolida, cornus, cosmos, deutzia, digitalis, erigeron, erodium, erysimum, erythronium, felicia, gazania, gypsophila, helenium, helianthemum, heliophila, hippeastrum, hyacinthus, hydrangea, iberis, ipomoea, ixia, jacaranda, ixia, kolkwitzia, lagerstroemia, lathyrus, lavatera, legousia, lewisia, linum, lobelia, lobularia, magnolia, malus, malva, mathiola, merendera, mimulus, myosotis, narcissus, nemesia, nicotiana, nopalxochitl, nymphaea, omphalodes, orthrosanthus, osteospermum, oxalis, paeonia, pelargonium, penstemon, pentas, pericallis, persicaria, platycodon, polemonium, polygala, potentilla, primula, prunus, puschkinia, rhododendron, rhodophlox, rose, saintpaulia, saponaria, saxifraga, scabiosa, schizostylis, schlumbergera, schilla, sedum, senecio, silene, solanum, spiraea, stachys, streptocarpus, syringa, tagetes, tanaecium, thun-

bergia, thymus, torenia, tropaeolum, verbenia, veronica, viburnum, vinca, viola, vitis, watsonia, and zinnia. The broad applicability of the modified *DFR* nucleic acids is based on the universal function of *DFR* in anthocyanin biosynthesis in divergent plant taxa. The parent plant used to practice the invention can be a wild type variant, a mutant which has been generated by the mutagenesis, or a transgenic line that has been generated by the recombinant techniques.

[0026] The invention also provides plant transformation vectors comprising at least a portion of substrate-specific *DFR* nucleic acids. In the preferred embodiments, the invention provides a plant transformation vector comprising at least a portion of DHK-specific *DFR* nucleic acids. Particularly preferred promoter to drive the expression of the DHK-specific *DFR* nucleic acids is the cauliflower mosaic virus 35S protein promoter. However, other constitutive promoters, tissue specific promoters, or inducible promoters can be also used.

[0027] The transformation of plants can be carried out in accordance with the invention by any of various transformation methods known to those skilled in the art of plant molecular biology. Particular methods for transformation include the transfer of nucleic acids into a plant cell by the microinjection, polyethylene glycol, electroporation, or microbombardment. Alternatively, plant cells can be transformed by *Agrobacterium* harboring vectors comprising at least a portion of modified *DFR* nucleic acids.

[0028] Regeneration of plants from the transformed cells can be carried out by any methods known to those skilled in the art. See, e.g., Methods in Enzymology, supra.; Methods in Enzymology, Vol 118; and Klee et al. Annual Review of Plant Physiology 38:467-486. Transformed cells or plants are selected based on their resistance to certain chemicals such as antibiotics or based on their phenotypes characterized by the increased content of pelargonidin-based pigments. The transformed plants can be self-fertilized or crossed with other plants. After the fertilization, the plants expressing at least portion of the modified *DFR* nucleic acids can be selected based on their resistance to certain chemicals such as antibiotics or based on their phenotypes characterized by the increased content of pelargonidin-based pigments. Alternatively, the transformed cells or a part of transformed plants can be grafted to other plants.

[0029] The following is presented as examples and is not to be construed as a limitation on the scope of the invention.

## EXAMPLE

### *Petunia* transformation

[0030] Leaf explants of the inbred *Petunia* W80 line (*an6*<sup>-</sup>, *ht1*<sup>-</sup>, *ht2*<sup>-</sup>, *h1*<sup>-</sup>, *ht2*<sup>-</sup>, *fr*<sup>-</sup>, and *rt*<sup>-</sup>) were transformed as described elsewhere except that leaf explants recently infected by *Agrobacterium tumefaciens* were rinsed with Murashige-Skoog solution containing 750 mg/L cefotaxime and then placed on media having 100 mg/L kanamycin sulfate and 500 mg/L cefotaxime (Johnson, et al. Plant J. 19:81-85 (1999)). Also, putative transformants were grown on MS media with vitamins, 30 g/L sucrose, 0.6% agar and 500 mg/L cefotaxime; after rooting the transformants were transferred to soil.

### Chimeric gene construction

[0031] Highly conserved regions of the *DFR* gene were identified by a multiple sequence alignment of a number of *DFRs*. The 5' region (*Gerbera DFR* portion) of each chimeric gene was synthesized from the *Gerbera DFR* cDNA clone using a primer containing the codon for the starting methionine of the *Gerbera DFR* gene (5'-GGC GAA AAT GGA AGA GGA TTC TCC-3') and a primer containing a conserved region of the *Gerbera DFR* gene (Chimera 1: 5'-AGC AGA TGA AGT GAA CAC TAG TTT CTT CAC-3'; Chimera 2: 5'-GGC TTT CTC TGC CAG AGT TTT TGA CAC GAA-3'; Chimera 3: 5'-GTG GGA CGA GCA AAT GTA TCT TCC TTT TGC-3'). The 3' region (*Petunia DFR* portion) of each chimeric gene was synthesized from the *Petunia DFRA* cDNA clone using a primer complementary to the three conserved regions (Chimera 1: 5'-TTC ACT TCA TCT GCT GGA ACT CTC GAT GTG; Chimera 2: 5'-CTG GCA GAG AAA GCC GCA ATG GAA GAA GCT-3'; Chimera 3: 5'-ATT TGT TCG TCC CAC CAT GCT ATC ATC TAC-3') and a primer containing the stop codon of the *Petunia DFRA* gene (5'-GCG CTA GAC TTC AAC ATT GCT TAA-3'). 5' and 3' regions were gel purified after PCR amplification. To assemble the full length chimeric gene the 5' and 3' region fragments were added to the same tube in roughly equal amounts and subjected to 25 PCR cycles (94°C 30", 55°C 30", 72°C 1:30). Full length chimeric genes (~1.1 kb) were purified from agarose gels. The chimeric genes were cloned into a vector containing the 35S CaMV promoter and NOS terminator. *Pfu* polymerase (Stratagene, La Jolla, CA) was used for all PCR reactions.

### Amino acid point mutant construction

[0032] *Gerbera DFR* genes containing one amino acid point mutation were made in a similar manner as the chimeric genes. The 5' region was synthesized using a primer having the *Gerbera DFR* starting methionine and a primer containing a single codon change. The 3' region was made with a complementary primer with the single codon change

and a primer having the stop codon of *Gerbera DFR*. The full length mutant sequence was assembled like the chimeric genes above. Each point mutant was cloned into a vector having the 35S CaMV promoter and NOS terminator. The mutagenized region of each mutant *DFR* was sequenced to ensure the correct residue was changed. Point mutants were then transformed into the W80 *Petunia* line. The transformants expressing the *DFR* genes were crossed with WR *Potunia* line (*dfrr*<sup>-</sup>, *F3'H*<sup>+/+</sup>) and WV *Petunia* line (*dfrr*<sup>-</sup>, *F3'5'H*<sup>+/+</sup>) to determine the substrate specificity of the mutated DFR. Mutations in other loci were not determined in these two *Petunia* lines.

#### TLC analysis

[0033] Anthocyanidins were separated on cellulose TLC plates as described (Johnson, et al. Plant J. 19:81-85 (1999)). Corollas were sometimes stored at 4°C for extended periods of time in methanol-0.5% HCl solution. Before adding iso-amylalcohol, the flower extracts were quantified at 530 nm to ensure uniform loading on the TLC plate. Anthocyanin standards were purchased from Apin Chemicals Ltd. (Oxfordshire, England).

#### Sequence alignment

[0034] Multiple sequence alignment of DFRs was done using ClustalW program.

SEQUENCE LISTING

<110> Korea Kumho Petrochemical Co., Ltd.

<120> Genetic sequences encoding substrate-specific  
dihydroflavonol 4- reductase and uses therefor

<130> Korea Kumho Petrochemical Co., Limited

<140>

<141>

<160> 32

<170> PatentIn Ver. 2.1

<210> 1

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic  
primer sequence

<400> 1

ggcgaaaatg gaagaggatt ctcc

24

<210> 2

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic  
primer sequence

<400> 2

agcagatgaa gtgaacacta gttttcttcac

30

<210> 3

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

# EP 1 182 257 A1

<223> Description of Artificial Sequence: synthetic  
primer sequence

<400> 3  
ggctttctct gccagagttt ttgacacgaa 30

<210> 4  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: synthetic  
primer sequence

<400> 4  
gtgggacgag caaatgtatc ttccttttgc 30

<210> 5  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: synthetic  
primer sequence

<400> 5  
ttcacttcat ctgctggaac tctcgatgtg 30

<210> 6  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: synthetic  
primer sequence

<400> 6  
ctggcagaga aagccgcaat ggaagaagct 30

<210> 7  
<211> 30



&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: synthetic  
primer sequence

&lt;400&gt; 7

atttgctcgt cccaccatgc tatcatctac

30

&lt;210&gt; 8

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: synthetic  
primer sequence

&lt;400&gt; 8

gcgctagact tcaacattgc ttaa

24

&lt;210&gt; 9

&lt;211&gt; 60

&lt;212&gt; PRT

&lt;213&gt; Gerbera sp.

&lt;400&gt; 9

Lys	Ala	Lys	Thr	Val	Lys	Lys	Leu	Val	Phe	Thr	Ser	Ser	Ala	Gly	Thr
1					5					10				15	

Val	Asn	Gly	Gln	Glu	Lys	Gln	Leu	His	Val	Tyr	Asp	Glu	Ser	His	Trp
	20						25						30		

Ser	Asp	Leu	Asp	Phe	Ile	Tyr	Ser	Lys	Lys	Met	Thr	Ala	Trp	Met	Tyr
	35						40					45			

Phe	Val	Ser	Lys	Thr	Leu	Ala	Glu	Lys	Ala	Ala	Trp
	50				55						60

&lt;210&gt; 10

&lt;211&gt; 60

&lt;212&gt; PRT

&lt;213&gt; Lilium sp.

EP 1 182 257 A1

<400> 10

Lys Ala Gly Thr Val Lys Arg Val Ile Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Val Asn Val Gln Glu Asn Gln Met Pro Glu Tyr Asp Glu Ser Ser Trp  
20 25 30

Ser Asp Val Asp Phe Cys Arg Arg Val Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Thr Leu Ala Glu Lys Ala Ala Trp  
50 55 60

<210> 11

<211> 60

<212> PRT

<213> Hordeum sp.

<400> 11

Glu Ala Gly Thr Val Lys Arg Ile Val Phe Thr Ser Ser Ala Gly Ser  
1 5 10 15

Val Asn Ile Glu Glu Arg Pro Arg Pro Ala Tyr Asp Gln Asp Asn Trp  
20 25 30

Ser Asp Ile Asp Tyr Cys Arg Arg Val Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Ala Leu Ala Glu Lys Ala Ala Met  
50 55 60

<210> 12

<211> 60

<212> PRT

<213> Antirrhinum sp.

<400> 12

Gln Ala Lys Thr Val Lys Lys Phe Ile Phe Thr Thr Ser Gly Gly Thr  
1 5 10 15

Val Asn Val Glu Glu His Gln Lys Pro Val Tyr Asp Glu Thr Asp Ser  
20 25 30

Ser Asp Met Asp Phe Ile Asn Ser Lys Lys Met Thr Gly Trp Met Tyr  
35 40 45

EP 1 182 257 A1

Phe Val Ser Lys Ile Leu Ala Glu Lys Ala Gly Met  
50 55 60

<210> 13

<211> 60

<212> PRT

<213> Petunia sp.

<400> 13

Lys Ala Asn Thr Val Lys Arg Leu Val Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Leu Asp Val Gln Glu Gln Gln Lys Leu Phe Tyr Asp Gln Thr Ser Trp  
20 25 30

Ser Asp Leu Asp Phe Ile Tyr Ala Lys Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Ala Ser Lys Ile Leu Ala Glu Lys Ala Ala Met  
50 55 60

<210> 14

<211> 60

<212> PRT

<213> Callistephus chinensis

<400> 14

Lys Ala Lys Thr Val Lys Lys Leu Val Tyr Thr Ser Ser Ala Gly Thr  
1 5 10 15

Val Asn Val Gln Glu Thr Gln Leu Pro Val Tyr Asp Glu Ser His Trp  
20 25 30

Ser Asp Leu Asp Phe Ile Tyr Ser Lys Lys Met Thr Ala Trp Met Tyr  
35 40 45

Phe Val Ser Lys Thr Leu Ala Glu Lys Ala Ala Met  
50 55 60

<210> 15

<211> 60

<212> PRT

<213> Daucus carota

<400> 15

EP 1 182 257 A1

Lys Ala Lys Thr Val Lys Lys Leu Ile Tyr Thr Ser Ser Ala Gly Thr  
1 5 10 15

Val Asn Val Arg Glu His Gln Leu Pro Val Tyr Asp Glu Ser Asn Trp  
20 25 30

Ser Asp Met Asp Phe Ile Tyr Ser Thr Lys Met Thr Ala Trp Met Tyr  
35 40 45

Phe Val Ser Lys Ser Leu Ala Glu Lys Ala Ala Trp  
50 55 60

<210> 16

<211> 60

<212> PRT

<213> Camellia japonica

<400> 16

Lys Ala Lys Thr Val Lys Arg Leu Val Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Val Asn Val Gln Glu His Gln Gln Pro Val Phe Asp Glu Asn Asn Trp  
20 25 30

Ser Asp Leu Asp Phe Ile Asn Lys Lys Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Thr Leu Ala Glu Lys Ala Ala Trp  
50 55 60

<210> 17

<211> 60

<212> PRT

<213> Arabidopsis sp.

<400> 17

Lys Ala Lys Thr Val Arg Arg Phe Val Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Val Asn Val Glu Glu His Gln Lys Asn Val Tyr Asp Glu Asn Asp Trp  
20 25 30

Ser Asp Leu Glu Phe Ile Met Ser Lys Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Ser Leu Ala Glu Lys Ala Ala Trp

EP 1 182 257 A1

50

55

60

<210> 18  
<211> 60  
<212> PRT  
<213> *Gentiana* sp.

<400> 18  
Lys Ala Lys Thr Val Lys Lys Leu Val Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Val Asp Val Gln Glu Gln Gln Lys Pro Val Tyr Asp Glu Asn Asp Trp  
20 25 30

Ser Asp Leu Asp Phe Ile Asn Ser Thr Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Ile Leu Ala Glu Lys Ala Ala Trp  
50 55 60

<210> 19  
<211> 60  
<212> PRT  
<213> *Ipomoea* sp.

<400> 19  
Lys Ala Lys Thr Val Lys Arg Leu Val Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Leu Asn Val Gln Pro Gln Gln Lys Pro Val Tyr Asp Glu Ser Cys Trp  
20 25 30

Ser Asp Leu Asp Phe Ile Tyr Ala Lys Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Ala Ser Lys Ile Leu Ala Glu Lys Glu Ala Trp  
50 55 60

<210> 20  
<211> 59  
<212> PRT  
<213> *Vitis* sp.

<400> 20  
Ala Lys Thr Val Arg Arg Leu Val Phe Thr Ser Ser Ala Gly Thr Val

EP 1 182 257 A1

1 5 10 15

Asn Ile Gln Glu His Gln Leu Pro Val Tyr Asp Glu Ser Cys Trp Ser  
20 25 30

Asp Met Glu Phe Cys Arg Ala Lys Lys Met Thr Ala Trp Met Tyr Phe  
35 40 45

Val Ser Lys Thr Leu Ala Glu Gln Ala Ala Trp  
50 55

<210> 21  
<211> 60  
<212> PRT  
<213> Forsynthia sp.

<400> 21  
Lys Ala Lys Thr Val Lys Arg Ile Val Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Val Asn Val Glu Glu His Gln Lys Ser Val Tyr Asp Glu Thr Asp Tyr  
20 25 30

Ser Asp Leu Asn Phe Ile Tyr Ser Lys Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Ile Leu Ala Glu Lys Val Ala Trp  
50 55 60

<210> 22  
<211> 60  
<212> PRT  
<213> Lycopersicon sp.

<400> 22  
Lys Ala Asn Thr Val Lys Arg Leu Val Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Leu Asp Val Gln Glu Asp Gln Lys Leu Phe Tyr Asp Glu Thr Ser Trp  
20 25 30

Ser Asp Leu Asp Phe Ile Tyr Ala Lys Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Ile Leu Ala Glu Lys Ala Ala Met  
50 55 60

EP 1 182 257 A1

<210> 23

<211> 60

<212> PRT

<213> Bromheadia sp.

<400> 23

Lys Ala Gly Ser Val Lys Arg Val Ile Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Val Asn Val Glu Glu His Gln Ala Ala Val Tyr Asp Glu Asn Ser Trp  
20 25 30

Ser Asp Leu His Phe Val Thr Arg Val Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Thr Leu Ala Glu Lys Ala Ala Trp  
50 55 60

<210> 24

<211> 60

<212> PRT

<213> Lotus sp.

<400> 24

Lys Ala Lys Thr Val Gln Arg Leu Val Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Leu Asn Ala Val Glu His Gln Lys Gln Met Tyr Asp Glu Ser Cys Trp  
20 25 30

Ser Asp Val Glu Phe Cys Arg Arg Val Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Thr Leu Ala Glu Gln Glu Ala Trp  
50 55 60

<210> 25

<211> 61

<212> PRT

<213> Rosa sp.

<400> 25

Lys Ala Lys Thr Val Arg Arg Leu Val Phe Thr Ser Ser Ala Gly Ser  
1 5 10 15

EP 1 182 257 A1

Val Asn Val Glu Glu Thr Gln Lys Pro Val Tyr Asn Glu Ser Asn Trp  
20 25 30

Ser Asp Val Glu Phe Cys Arg Arg Val Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Ala Ser Lys Thr Leu Ala Glu Glu Gln Glu Ala Trp  
50 55 60

<210> 26  
<211> 60  
<212> PRT  
<213> Glycine sp.

<400> 26  
Lys Ala Lys Thr Val Arg Arg Leu Ile Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Leu Asn Val Ile Glu Arg Gln Lys Pro Val Phe Asp Asp Thr Cys Trp  
20 25 30

Ser Asp Val Glu Phe Cys Arg Arg Val Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Thr Leu Ala Glu Lys Glu Ala Trp  
50 55 60

<210> 27  
<211> 59  
<212> PRT  
<213> Zea sp.

<400> 27  
Glu Ala Gly Thr Val Arg Arg Ile Val Phe Thr Ser Ser Ala Gly Thr  
1 5 10 15

Val Asn Leu Glu Glu Arg Gln Arg Pro Val Tyr Asp Glu Glu Ser Trp  
20 25 30

Thr Asp Val Asp Phe Cys Arg Arg Val Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Thr Leu Ala Glu Lys Ala Ala  
50 55



<210> 28  
 <211> 59  
 <212> PRT  
 <213> Sorghum sp.

<400> 28  
 Glu Ala Gly Thr Val Arg Arg Ile Val Phe Thr Ser Ser Ala Gly Thr  
           1                  5                  10                  15  
 Val Asn Ile Glu Glu Arg Gln Arg Pro Val Tyr Asp Gln Asp Asn Trp  
                   20                  25                  30  
 Ser Asp Val Asp Phe Cys Gln Arg Val Lys Met Thr Gly Trp Met Tyr  
                   35                  40                  45  
 Phe Val Ser Lys Ser Leu Ala Glu Lys Ala Ala  
           50                  55

<210> 29  
 <211> 60  
 <212> PRT  
 <213> Medicago sp.

<400> 29  
 Lys Ala Lys Thr Val Arg Arg Leu Ile Tyr Thr Ser Ser Ala Gly Thr  
           1                  5                  10                  15  
 Leu Asn Val Thr Glu Asp Gln Lys Pro Leu Trp Asp Glu Ser Cys Trp  
                   20                  25                  30  
 Ser Asp Val Glu Phe Cys Arg Arg Val Lys Met Thr Gly Trp Met Tyr  
                   35                  40                  45  
 Phe Val Ser Lys Thr Leu Ala Glu Gln Glu Ala Trp  
           50                  55                  60

<210> 30  
 <211> 59  
 <212> PRT  
 <213> Oryza sp.

<400> 30  
 Ala Gly Thr Val Lys Arg Ile Val Phe Thr Ser Ser Ala Gly Thr Val  
           1                  5                  10                  15

EP 1 182 257 A1

Asn Ile Glu Glu Arg Gln Arg Pro Ser Tyr Asp His Asp Asp Trp Ser  
20 25 30

Asp Ile Asp Phe Cys Arg Arg Val Lys Met Thr Gly Trp Met Tyr Phe  
35 40 45

Val Ser Lys Ser Leu Ala Glu Lys Ala Ala Met  
50 55

<210> 31

<211> 60

<212> PRT

<213> Fragaria sp.

<400> 31

Lys Ala Lys Thr Val Arg Arg Leu Val Phe Thr Ser Ser Ala Gly Ala  
1 5 10 15

Val Ala Ile Glu Glu His Pro Lys Glu Val Tyr Ser Glu Asn Asn Trp  
20 25 30

Ser Asp Val Val Phe Cys Arg Lys Val Lys Met Thr Gly Trp Met Tyr  
35 40 45

Phe Val Ser Lys Thr Leu Ala Glu Gln Ala Ala Trp  
50 55 60

<210> 32

<211> 32

<212> PRT

<213> Gerbera sp.

<400> 32

Phe Thr Ser Ser Ala Gly Thr Val Asn Gly Gln Glu Lys Gln Leu His  
1 5 10 15

Val Tyr Asp Glu Ser His Trp Ser Asp Leu Asp Phe Ile Tyr Ser Lys  
20 25 30

## Claims

1. The substrate specific *DFR* nucleic acids and encoding DFRs that have altered amino acid sequences at the substrate specificity determining region due to the altered nucleic acid sequences encoding amino acids at the substrate specificity determining region.
2. The substrate specific *DFR* nucleic acids and encoding DFR of claim 1 which catalyzes DHK preferentially over DHQ and DHM.
3. The substrate specific *DFR* nucleic acids and encoding DFR of claim 2 which has altered amino acid at 134<sup>th</sup> residue of *Gerbera* DFR or the corresponding amino acid residues of DFRs from other species.
4. The substrate specific *DFR* nucleic acids and encoding DFR of claim 3 which has leucine instead of asparagine at the said residue.
5. An angiosperm plant comprising at least one cell transformed with a vector comprising at least a portion of the substrate specific *DFR* nucleic acids of claim 1 and wherein said plant has the increased content of one class of pigment preferentially.
6. An angiosperm plant comprising at least one cell transformed with a vector comprising at least a portion of the substrate specific *DFR* nucleic acids of claim 2 and wherein said plant has the increased content of pelargonidin-based pigments.
7. An angiosperm plant comprising at least one cell transformed with a vector comprising at least a portion of the substrate specific *DFR* nucleic acids of claim 3 and wherein said plant has the increased content of pelargonidin-based pigments.
8. An angiosperm plant comprising at least one cell transformed with a vector comprising at least a portion of the substrate specific *DFR* nucleic acids of claim 4 and wherein said plant has the increased content of pelargonidin-based pigments.
9. The plant of claim 5 wherein said vector comprises a promoter operably linked to said *DFR* nucleic acids.
10. The plant of claim 9 wherein said promoter comprises a constitutive promoter.
11. The plant of claim 10 wherein said promoter comprises a cauliflower mosaic virus promoter.
12. The plant of claim 9 wherein said promoter comprises a tissue specific promoter.
13. The plant of claim 9 wherein said promoter comprises an inducible promoter.
14. A vector capable of transforming a plant cell to increase the content of one class of pigments preferentially in a plant containing said cell, said vector comprising at least a portion of the substrate specific *DFR* nucleic acids operably linked to a promoter.
15. The vector of claim 14 wherein said substrate specific *DFR* nucleic acids are DHK-specific *DFR* nucleic acids.
16. The vector of claim 14 wherein said one class of pigments is pelargonidin-based pigments.
17. The vector of claim 14 wherein said promoter is cauliflower mosaic virus promoter.
18. A method for producing a plant having a phenotype **characterized by** the increased content of one class of pigments, said method comprising at least a step of:
  - transforming plant cells with a vector comprising at least a portion of substrate specific *DFR* nucleic acids operably linked to a promoter;
  - regenerating plants from one or more of the transformed plant cells; and selecting at least one plant having said phenotype.

**EP 1 182 257 A1**

19. The method of claim 18 wherein said one class of pigments is pelargonidin-based pigments.
20. The method of claim 18 wherein said promoter comprises a cauliflower mosaic virus promoter.
21. An angiosperm plant produced according to the method of claim 18.

FIG. 1A




FIG. 1B



FIG. 1C




FIG. 2

```

Gerbera      : KAKTVKKLVFTSSAGTVWVGQEKQLHVYDESHWSLDLFIYSKKMTAMMYFVSKTLAEKAAW
Lilium       : KAGTVKRVIFTSSAGTVWVQENQNPPEYDESNWSVDVFCRRVKMTGMYFVSKTLAEKAAW
Hordeum      : EAGTVKRVIFTSSAGTVWVIEERPRPAYDQDNWSDIDYCRVKMTGMYFVSKTLAEKAAW
Antirrhinum  : QAKTVKKPIFTSSAGTVWVVEHQKPVYDETSDDSDIFNSKKMTGMYFVSKTLAEKAAW
Petunia      : KANTVKKLVFTSSAGTLDVQEQQKLFYDQTSWSDLDFIYAKKMTGMYFVSKTLAEKAAW
Callistephus: KAKTVKKLVFTSSAGTVWVQETQLPVYDESHWSLDLFIYSKKMTAMMYFVSKTLAEKAAW
Daucus       : KAKTVKKLIYFTSSAGTVWVREHQLPVYDESNWSDDHDFIYTKMTAMMYFVSKTLAEKAAW
Camellia     : KAKTVKKLVFTSSAGTVWVQEQHQPVDYDENWSDLDFINSKKMTGMYFVSKTLAEKAAW
Arabidopsis  : KAKTVRRPVFTSSAGTVWVVEHQKPVYDENWSDLEFIMSKKMTGMYFVSKTLAEKAAW
Gentiana     : KAKTVKKLVFTSSAGTVDVQEQKPVYDENWSDLDFINSTKMTGMYFVSKTLAEKAAW
Ipomoea      : KAKTVKKLVFTSSAGTLDVQEQKPVYDESNWSLDLFIYAKKMTGMYFVSKTLAEKAAW
Vitis        : AKTVKKLVFTSSAGTVWVQEQHQLPVYDESNWSDDHDFIYAKKMTGMYFVSKTLAEKAAW
Forsythia    : KAKTVKRVIFTSSAGTVWVVEHQKSVYDETDYSDLNF IYAKKMTGMYFVSKTLAEKAAW
Lycopersicon: KANTVKKLVFTSSAGTLDVQEQKLFYDETSWSDLDFIYAKKMTGMYFVSKTLAEKAAW
Bromheadia   : KAGSVKRVIFTSSAGTVWVVEHQAAVYDENWSDLHFVTRVKMTGMYFVSKTLAEKAAW
Lotus        : KAKTVKKLVFTSSAGTLDVQEQHQPVDYDENWSDLDFINSKKMTGMYFVSKTLAEKAAW
Rosa         : KAKTVRRPVFTSSAGTVWVVEHQKPVYDENWSDLEFIMSKKMTGMYFVSKTLAEKAAW
Glycine      : KAKTVRRLIYFTSSAGTLDVQEQKPVYDETDYSDLNF IYAKKMTGMYFVSKTLAEKAAW
Zea          : EAGTVRRIVFTSSAGTVWVVEHQKPVYDESNWSDVDFCRRVKMTGMYFVSKTLAEKAAW
Sorghum      : EAGTVRRIVFTSSAGTVWVVEHQKPVYDESNWSDVDFCRRVKMTGMYFVSKTLAEKAAW
Medicago    : KAKTVRRLIYFTSSAGTLDVQEQHQPVDYDENWSDLDFINSKKMTGMYFVSKTLAEKAAW
Oryza        : AGTVKRVIFTSSAGTVWVVEHQKPVYDESNWSDIDYCRVKMTGMYFVSKTLAEKAAW
Fragaria     : KAKTVRRPVFTSSAGTVWVVEHQKPVYDESNWSDVDFCRRVKMTGMYFVSKTLAEKAAW

```

FIG. 3A



FIG. 3B



FIG. 3C




FIG. 4A





FIG. 4B







European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 00 30 8002

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Inv.C1.7)
X	JOHNSON, E.T. ET AL.: "Cymbidium hybrida dihydroflavonol 4-reductase does not efficiently reduce dihydrokaempferol to produce orange pelargonidin-type anthocyanins" PLANT J., vol. 19, no. 1, 1999, pages 81-85, XP002173168 * figure 3B * * page 84, left-hand column, paragraph 1 - paragraph 3 *	1-21	C12N15/53 C12N9/02 C12N9/04 C12N15/82 A01H5/00
X	EP D 316 797 A (MAX PLANCK GESELLSCHAFT) 24 May 1989 (1989-05-24) * column 2, paragraph 3 *	18-21	
			TECHNICAL FIELDS SEARCHED (Inv.C1.7)
			C12N A01H
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 25 July 2001	Examiner Mata Vicente, T.
CATEGORY OF CITED DOCUMENTS			
<p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>* : member of the same parent family, corresponding document</p>			

PRO-FORMA (ISO 10305-2:1992)

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 30 8002

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

25-07-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0316797 A	24-05-1989	DE 3738657 C	18-05-1989
		DE 3885061 D	25-11-1993
		ES 2059472 T	16-11-1994
		IL 88307 A	18-08-1993
		JP 2002305 A	08-01-1990
		JP 2996663 B	11-01-2000
		US 5410096 A	25-04-1995
-----			

EPC/EPN/PA/48

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82